

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Exhibit "E"

CENTER FOR DRUG EVALUATION AND RESEARCH

Application Number: NDA 20062/S027

APPROVAL LETTER



DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service

Food and Drug Administration
Rockville MD 20857

AUG 24 1999

DR

NDA 20-062/S-027

Marion Merrell Dow (Europe) AG
as General Partner of
Carderm Capital L.P.
c/o Westbroke Limited
Attention: Mr. Carlos A. Austin
Richmond House
12 Par-la Ville Road
P.O. Box HM 1022
Hamilton HM DX
Bermuda

Dear Mr. Austin:

Please refer to your supplemental new drug application dated January 7, 1999, received January 11, 1999, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Cardizem CD (diltiazem hydrochloride) 180, 240, 300 and 360 mg Capsules.

We acknowledge receipt of your submissions dated May 11, June 18, and July 27, 1999. Your submission of June 18, 1999 constituted a complete response to our May 7, 1999 action letter.

This supplemental new drug application provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules. Final printed labeling has been revised to incorporate information regarding this new dosage strength. In addition, the How Supplied statement was revised to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

We have completed the review of this supplemental application, as amended, and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the submitted final printed labeling (package insert and immediate container and carton label submission dated June 18, 1999). Accordingly, the supplemental application is approved effective on the date of this letter.

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

NDA 20-062/S-027
Page 2

If you have any questions, please contact:

David Roeder
Regulatory Health Project Manager
(301) 594-5313

Sincerely,

RSI 8/22/21

Raymond J. Lipicky, M.D.
Director
Division of Cardio-Renal Drug Products
Office of Drug Evaluation I
Center for Drug Evaluation and Research

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

APPROVABLE LETTER



DEPARTMENT OF HEALTH & HUMAN SERVICES

Food and Drug Administration
Rockville MD 20857

NDA 20-062/S-027

MAY -7 1999

Marion Merrell Dow (Europe) AG
as General Partner of
Cardem Capital L.P.
c/o Westbroke Limited
Attention: Carlos A. Austin
Richmond House
12 Par-la Ville Road
P.O. Box HM 1022
Hamilton HM DX
Bermuda

Dear Mr. Austin:

Please refer to your supplemental new drug application dated January 7, 1999, received January 11, 1999, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Cardizem CD (diltiazem HCl) 120, 180, 240 and 300 mg Capsules.

We acknowledge receipt of your submission dated March 5, 1999.

This supplement provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules.

We have completed the review of this application, as amended, and it is approvable. Before this application may be approved, however, it will be necessary for you to submit final printed labeling (FPL) revised as follows:

The Storage Statement should be revised in the package insert and the container labels to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP
Controlled Room Temperature].

In addition, all previous revisions as reflected in the most recently approved labeling must be included. To facilitate review of your submission, please provide a highlighted or marked-up copy that shows the changes that are being made.

Please note that stability data at the 12-month time point for the 360 mg strength capsule at 25°C/60% RH and at 30°C/60% RH should be submitted in support of a 24-month expiration date.

Validation of the regulatory methods has not been completed. At the present time, it is the policy of the Center not to withhold approval because the methods are being validated. Nevertheless, we expect your continued cooperation to resolve any problems that may be identified.

Please submit 20 copies of the final printed labeling, ten of which are individually mounted on heavy weight paper or similar material.

If additional information relating to the safety or effectiveness of this drug becomes available, revision of the labeling may be required.

Within 10 days after the date of this letter, you are required to amend the supplemental application, notify us of your intent to file an amendment, or follow one of your other options under 21 CFR 314.110. In the absence of any such action FDA may proceed to withdraw the application. Any amendment should respond to all the deficiencies listed. We will not process a partial reply as a major amendment nor will the review clock be reactivated until all deficiencies have been addressed.

This product may be considered to be misbranded under the Federal Food, Drug, and Cosmetic Act if it is marketed with this change prior to approval of this supplemental application.

If you have any questions, please contact:

Mr. David Roeder
Regulatory Health Project Manager
(301) 594-5313

Sincerely yours,

151 5/7/69
Raymond J. Lipicky, M.D.
Director
Division of Cardio-Renal Drug Products
Office of Drug Evaluation I
Center for Drug Evaluation and Research

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027


FINAL PRINTED LABELING

Reviewed by: A. S. L. 8-29-88

APPROVED

Prescribing information as of May 1999
CARDIZEM® CD
 (diltiazem HCl)
 Capsules

CALAZEM® (difenhydramide) is a calcium ion influx inhibitor (slow channel blocker or calcium antagonist). Chemically, difenhydramide is 15-benzocyclohexyl-4(5H)-one-3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2(1-methylpiperidin-4-yl)-1H-imidazole-1-carboxylate. The chemical structure is:



Dilantin hydrochloride is a white to off-white crystalline powder with a bitter taste. It is soluble in water, slightly soluble in alcohol. Its molecular weight is 356.29.

CAPAZOLIN CD is formulated as once-a-day extended release capsules containing either 120 mg, 180 mg, 240 mg, 360 mg, or 360 mg dilantin hydrochloride. The 120 mg, 180 mg, 240 mg, and 360 mg capsules also contain: black iron oxide, polyethylene glycol, PEG 350, fumaric acid, croscarmellose sodium, starch, lactose, polydioxanone, talc, wax, and other inactive ingredients. Each 360 mg capsule also contains: hard paraffin, polyethylene glycol, PEG 350, polydioxanone, K17, calcium lauryl sulfate, stearic acid, sucrose, talc, titanium dioxide, and other inactive ingredients.

CLINICAL PHARMACOLOGY
The therapeutic effects of CAPROXEN® CO are believed to be related to its ability to inhibit the action of calcium ions during membrane depolarization of cardiac and vascular smooth muscle.

Heartburners of Japan
Aspirin. CARDON CO produces the antipyrretic analgesic primarily by synthesis of acetanilide, salicylic acid and the resulting aspirin. The synthesis of blood thinners is a function of the degree of reduction of the aspirin. Aspirin is a blood thinner, antipyrretic and analgesic. There is only a modest risk in blood coagulation in aspirin therapy.

Aspirin. CARDON CO has been shown to produce increases in blood flow, blood viscosity due to its ability to reduce blood viscosity. This is accompanied by a decrease in blood viscosity. This is accompanied by a decrease in blood viscosity. This is accompanied by a decrease in blood viscosity.

2. **Ischemic models.** Ischemic infarction is the most common epidemiologic cause of stroke. Ischemic infarction is characterized by a focal area of brain tissue that has died due to a lack of blood flow. The most common cause of ischemic infarction is a blood clot that has formed in the carotid artery, which supplies the brain with blood. The clot can break off and travel to the brain, where it can block the artery and cause a stroke. Other causes of ischemic infarction include atherosclerosis, which is a buildup of plaque in the arteries, and a blood clot that has formed in the heart and traveled to the brain.

Like other calcium channel antagonists, diltiazem decreases sinus and sinoatrial conduction in isolated tissues and has a negative inotropic effect in isolated preparations. In the intact animal, prolongation of the AH interval can be seen at further doses.

In man, different pressures operate on and expunge through the coronary artery system. It causes a rise in peripheral vascular resistance and increases blood pressure in aortic and coronary arteries, and in coronary resistance studies in the dog, the technique heart disease, increases the peripheral vascular resistance for any given work load. Studies to date, primarily in patients with good ventricular function, have not shown a significant negative inotropic effect; however, a negative inotropic effect, and left ventricular dysfunction, and left ventricular failure, if the pressure has not been affected, has been observed in patients with severe left ventricular failure. Patients with poor ventricular function, and increased left ventricular pressure, have been shown to have a pronounced impairment of ventricular function. There are as yet no data on the interaction of diuretics and beta-blockers in patients with poor ventricular function. There is some data to suggest slightly reduced by diuretics.

CHLORAZEPATE (C) lowered cardiac diastolic blood pressure in an dose-dependent manner over the 24-hr period. The maximum change in diastolic blood pressure occurred at 12 hr, for placebo, 100 mg, 200 mg, and 400 mg were -2.0 , -3.5 , -5.5 , and -6.5 mm Hg, respectively. Postural hypotension is infrequently noted with C, especially assuming an upright posture. C is not associated with the chronic antihypertensive effects of thiazide diuretics. CHLORAZEPATE (C) decreases myocardial contractility (stroke volume), and produces a slight decrease or no change in heart rate. The dynamic response, measured as the rate of pressure rise in the left ventricle, is also decreased. Myocardial contractility is usually reduced. CHLORAZEPATE (C) produces changes in peripheral vascular resistance. It increases arterial resistance, but decreases peripheral vascular resistance. CHLORAZEPATE (C) reduces the renal-angiotensin system. CHLORAZEPATE (C) reduces the effects of angiotensin II.

double-blind, parallel, dose-response study of doses from 60 to 90 mg once daily. CHOLEZEN could lead the way to investigation of

CHS in a later summer over the 5 dose groups studied. The observed time to termination of CHS utilizing a three-minute assessment of trough, for groups 0, 50, 120, 240, 480, and 960 mg was 21, 43, 50, 50, and 68 days, respectively. Because of CARDEX CO were decreased, overall mean frequency of CHS once daily, or plasma concentration in a 12-hour study patients receiving cacosulfate tablets with long-acting salicylate or beta-blocker. A significant increase in time to termination of

and a significant decrease in diastolic frequency was noted. In this trial the overall rate of adverse events in the ZEN CO treatment group was low as the placebo group. No major differences in doses of digoxin or AH conduction time were observed and efficacy was equally marked by supraventricular tachycardia. The results of a study involving 10 patients and of 200 mg of CaH_2Zn in six patients, the average conduction time was 14% with a decrease of greater than 50% in AH block. Differences noted in the conduction of the AH interval is pronounced in patients with severe heart block. In patients with slow conduction, differences in the AH interval are noted. The AH interval is noted to be

3 These are administration of CAROZEM to patients in doses of up to 540 mg/day has resulted in small increases in PR interval, and on occasion produced abnormal prolongation. (See WARNINGS.)

Pharmacokinetics and Bioeffects
Chromium is well absorbed from the gastrointestinal tract and is subject to an extensive first-pass effect, giving an absolute bioavailability (compared to intravenous administration) of about 10%. CARCOTE[®] undergoes extensive metabolism, with only 2% to 4% of the injected dose appearing in the urine. Drugs which induce or inhibit hepatic metabolism may alter chromium disposition.

Toxicity
Toxic radioactivity measurements following short IV administrations in healthy volunteers suggests the presence of other substances in the preparation, which allow higher concentrations than those of chromium and are more slowly eliminated; half-life of radioactivity in about 20 hours compared to 2 to 5 hours for chromium.

by very binding studies show CARPOZINE is 70% to 80% bound to plasma proteins. Competitive *in vitro* studies have shown that plasma protein binding of CARPOZINE is not significantly affected by therapeutic concentrations of digoxin, hydrochlorothiazide, furosemide, or propranolol. CARPOZINE is not metabolized by the liver and is excreted in the urine as the parent drug. Following intravenous administration, the elimination half-life is 1.5 hours. In multiple drug administration, the elimination half-life is 1.5 hours. (See table 1.)

Plasma concentrations of 3.0 to 10.0 mg/L are present in the plasma of healthy adults. About 25% of the parent drug is excreted in the urine as the parent drug. About 75% is excreted as the primary metabolite, 2-hydroxy-carpozine.

When therapeutic plasma concentrations appear to be in the range of 3.0 to 10.0 mg/L, there is a decrease in plasma protein binding as the plasma concentrations are increased. The plasma albumin is slightly increased with increasing plasma concentrations. The plasma albumin is decreased with increasing plasma concentrations. The plasma albumin is decreased with increasing plasma concentrations. The plasma albumin is decreased with increasing plasma concentrations.

When patients with chronic renal insufficiency are treated with CARPOZINE, the elimination half-life is increased. The elimination half-life is increased in patients with chronic renal insufficiency. The elimination half-life is increased in patients with chronic renal insufficiency. The elimination half-life is increased in patients with chronic renal insufficiency.

A study in patients with severe renal insufficiency showed that the elimination half-life of CARPOZINE is increased. The elimination half-life is increased in patients with severe renal insufficiency. The elimination half-life is increased in patients with severe renal insufficiency. The elimination half-life is increased in patients with severe renal insufficiency.

CARBOGEN CO Capentec, When reacted to a mixture of CARBOGEN (a mixture of ethylene oxide, more than 90% ethylene oxide) and water, the reaction is observed from the first day of the reaction. A single drop of water is sufficient to initiate the reaction. The reaction is exothermic and produces heat within seconds and much plasma forms. The reaction is self-sustaining and does not require the addition of any other chemicals. The reaction is exothermic and produces heat within seconds and much plasma forms. The reaction is self-sustaining and does not require the addition of any other chemicals.

CAUTIONS AND WARNINGS
ACEEM CO is indicated for the treatment of hypertension. It may be used alone or in combination with antihypertensive medications.

ACEEM CO is indicated for the treatment of chronic stable angina pectoris due to coronary artery

21M is compromised in (1) with sick sinus syndrome in the presence of a rising ventricular pacemaker, (2) patients with hypotension (BP less than 90/60 mm Hg), (3) patients with severe pulmonary edema, (4) patients with severe renal insufficiency, and (5) patients with acute and/or chronic pulmonary embolism. The procedure was discontinued in 10 patients because of the above-mentioned contraindications.

the Conduction. CAROTID sinus AV node activity may be slowed significantly during sinus tachycardia except in patients with sick sinus syndrome. This effect may result in obscuring slow rates (particularly in patients with sinus dysrhythmia) or 2:1 or third-degree AV block (22% patients, or 0.4%).

Combined use of diltiazem with digoxin or dopamine may enhance effects on cardiac output. A patient with ventricular angina developed a 2° AV block (2 to 3 sec) after a single dose of diltiazem. (See ADVERSE

2. **Coagulative Heart Failure**
Although diltiazem has a negative inotropic effect in isolated animal tissue preparations, hemodynamic studies in humans with normal ventricular function have shown that

fraction have not shown a reduction in cardiac output or consistent negative effects on contractility (p/pv). An early study of oral disodium in patients with impaired ventricular function (fraction 24%, 6%) showed improvement in indices of ventricular function without significant decrease in contractile function (dp/dt). Worsening of congestive heart failure has been reported in patients with pre-existing impairment of ventricular function. Experience with the use of disodium (amilofrudine hydrochloride) in combination with beta-blockers in patients with impaired ventricular function is limited. Caution should be exercised when using this combination.

4. **Acute Hepatic Injury.** Mild elevations of transaminases with and without concomitant elevation in alkaline phosphatase and bilirubin have been observed in clinical studies. Such elevations were usually transient and frequently resolved even with continued diazepam treatment. In rare instances, significant elevations in enzymes such as alkaline phosphatase, LDH, SGOT, SGPT, and other phenomena suggestive of acute hepatic injury have been observed. These reactions tended to occur early after therapy initiation (1 to 8 weeks) and have been reversible upon discontinuation of drug therapy. The relationship to CARBODOL is uncertain in some cases, but probable in some. (See

PRELIMINARY

CARBAMID (alkaloids hydrochloride) is continuously autoabsorbed by the lungs and excreted by the kidneys and liver. As with aspirin, the laboratory parameters of renal and hepatic function should be monitored at regular intervals. The drug should be used with caution in patients with impaired renal or hepatic function, in patients who are on long-term therapy and in patients who are on therapy with drugs which may cause changes in clearance with hepatic damage. In special cases of renal disease, doses of 125 mg/kg and below, and doses of 100 mg/kg and below, respectively, should be used. In patients with hepatic changes, doses of 20 mg/kg were associated with hepatic changes, whereas, these changes were not observed with doses of 10 mg/kg. In patients with renal disease, doses of 20 mg/kg were associated with hepatic changes. In patients with hepatic changes, doses of 20 mg/kg were associated with hepatic changes. In patients with hepatic changes, doses of 20 mg/kg were associated with hepatic changes.

to the potential for additive and synergistic effects. CAROZIN is indicated for treatment of patients with angina in patients receiving CAROZIN concomitantly with other drugs known to affect cardiac conduction. The following drugs are known to affect cardiac conduction: (1) Class I antiarrhythmic drugs (e.g., quinidine, procainamide, flecainide, etc.) which may have an additive effect with CAROZIN in causing low QRS conduction (2) Class II antiarrhythmic drugs (e.g., propranolol, etc.) which may enhance the negative inotropic effect of CAROZIN. (See CONTRAINDICATIONS.)

As a precaution, care should be exercised when treating patients with multiple medications. CAROZIN undergoes biotransformation with many drugs, and may interact with other drugs. In addition, CAROZIN may alter effects which the same results of biotransformation. In addition, the effects of CAROZIN may be altered by the effects of metabolism. Especially, patients with renal and/or hepatic impairment, a decrease of biotransformation, and/or a decrease of excretion may experience a decrease in the therapeutic effect. In addition, patients with hepatic impairment may experience a decrease in the therapeutic effect. In addition, patients with hepatic impairment may experience a decrease in the therapeutic effect.

ble. Controlled and uncontrolled domestic studies suggest an excellent way of CAPOZEN auto-injection is usually used. But available data are not used to predict the effects of current treatment in patients at vascular dysfunction or conduction abnormalities. Treatment of CAPOZEN (disodium phosphate) consistently associated in two normal volunteers to increased venous pressure in all subjects and stability of propofol used and approximately 50% in

One of the most common reasons for the decision to use a biologic is the physician's or veterinarian's therapy is reduced or withdrawn in compliance with prescriptions or regulations. The veterinarian's decision may be forced. (See *Wetters*.)

Another study in the healthy volunteers has shown that a significant increase in peak diastolic blood pressure (24%) and area-under-the-curve (25%) after a 1-week course of treatment of 1200 mg per day and a single dose of 2400 mg to 2800 mg. The drug produced greater, more prolonged, and more potent effects than the placebo. The effect may be mediated by direct or indirect inhibition of renin-angiotensin conversion of angiotensinogen to angiotensin II. The enzyme system responsible for the first-step metabolism of renin. Patients currently receiving antihypertensive therapy should be carefully monitored for changes in pharmacological effect, including edema and congestive heart failure. Concomitant therapy with diuretics. As adjustments in the dosage may be warranted.

Labeling: ORIGINAL
FDA No: 20-168 Re'd. 6-21-99
Reviewed by: R. Smith 8-24-99

DATE: 6/18/99	CICERO JOB NO.: 13571
PRODUCT: LABEL, CARDIZEM CD, CAPSULES 360 MG, 90 CT.	
PRINTER: NEW JERSEY PACKAGING	
FIG 201	FIG 202

APPROVALS

FUNCTIONAL

PROOFREADING

REGULATORY

APPROVED
AUG 24 1999


⊕ I PMS 200 PMS 201 PMS 202 PMS 203 PMS 204 PMS 205 PMS 206 PMS 207 PMS 208 PMS 209 PMS 210 PMS 211 PMS 212 PMS 213 PMS 214 PMS 215 PMS 216 PMS 217 PMS 218 PMS 219 PMS 220 PMS 221 PMS 222 PMS 223 PMS 224 PMS 225 PMS 226 PMS 227 PMS 228 PMS 229 PMS 230 PMS 231 PMS 232 PMS 233 PMS 234 PMS 235 PMS 236 PMS 237 PMS 238 PMS 239 PMS 240 PMS 241 PMS 242 PMS 243 PMS 244 PMS 245 PMS 246 PMS 247 PMS 248 PMS 249 PMS 250 PMS 251 PMS 252 PMS 253 PMS 254 PMS 255 PMS 256 PMS 257 PMS 258 PMS 259 PMS 260 PMS 261 PMS 262 PMS 263 PMS 264 PMS 265 PMS 266 PMS 267 PMS 268 PMS 269 PMS 270 PMS 271 PMS 272 PMS 273 PMS 274 PMS 275 PMS 276 PMS 277 PMS 278 PMS 279 PMS 280 PMS 281 PMS 282 PMS 283 PMS 284 PMS 285 PMS 286 PMS 287 PMS 288 PMS 289 PMS 290 PMS 291 PMS 292 PMS 293 PMS 294 PMS 295 PMS 296 PMS 297 PMS 298 PMS 299 PMS 300 PMS 301 PMS 302 PMS 303 PMS 304 PMS 305 PMS 306 PMS 307 PMS 308 PMS 309 PMS 310 PMS 311 PMS 312 PMS 313 PMS 314 PMS 315 PMS 316 PMS 317 PMS 318 PMS 319 PMS 320 PMS 321 PMS 322 PMS 323 PMS 324 PMS 325 PMS 326 PMS 327 PMS 328 PMS 329 PMS 330 PMS 331 PMS 332 PMS 333 PMS 334 PMS 335 PMS 336 PMS 337 PMS 338 PMS 339 PMS 340 PMS 341 PMS 342 PMS 343 PMS 344 PMS 345 PMS 346 PMS 347 PMS 348 PMS 349 PMS 350 PMS 351 PMS 352 PMS 353 PMS 354 PMS 355 PMS 356 PMS 357 PMS 358 PMS 359 PMS 360 PMS 361 PMS 362 PMS 363 PMS 364 PMS 365 PMS 366 PMS 367 PMS 368 PMS 369 PMS 370 PMS 371 PMS 372 PMS 373 PMS 374 PMS 375 PMS 376 PMS 377 PMS 378 PMS 379 PMS 380 PMS 381 PMS 382 PMS 383 PMS 384 PMS 385 PMS 386 PMS 387 PMS 388 PMS 389 PMS 390 PMS 391 PMS 392 PMS 393 PMS 394 PMS 395 PMS 396 PMS 397 PMS 398 PMS 399 PMS 400 PMS 401 PMS 402 PMS 403 PMS 404 PMS 405 PMS 406 PMS 407 PMS 408 PMS 409 PMS 410 PMS 411 PMS 412 PMS 413 PMS 414 PMS 415 PMS 416 PMS 417 PMS 418 PMS 419 PMS 420 PMS 421 PMS 422 PMS 423 PMS 424 PMS 425 PMS 426 PMS 427 PMS 428 PMS 429 PMS 430 PMS 431 PMS 432 PMS 433 PMS 434 PMS 435 PMS 436 PMS 437 PMS 438 PMS 439 PMS 440 PMS 441 PMS 442 PMS 443 PMS 444 PMS 445 PMS 446 PMS 447 PMS 448 PMS 449 PMS 450 PMS 451 PMS 452 PMS 453 PMS 454 PMS 455 PMS 456 PMS 457 PMS 458 PMS 459 PMS 460 PMS 461 PMS 462 PMS 463 PMS 464 PMS 465 PMS 466 PMS 467 PMS 468 PMS 469 PMS 470 PMS 471 PMS 472 PMS 473 PMS 474 PMS 475 PMS 476 PMS 477 PMS 478 PMS 479 PMS 480 PMS 481 PMS 482 PMS 483 PMS 484 PMS 485 PMS 486 PMS 487 PMS 488 PMS 489 PMS 490 PMS 491 PMS 492 PMS 493 PMS 494 PMS 495 PMS 496 PMS 497 PMS 498 PMS 499 PMS 500 PMS 501 PMS 502 PMS 503 PMS 504 PMS 505 PMS 506 PMS 507 PMS 508 PMS 509 PMS 510 PMS 511 PMS 512 PMS 513 PMS 514 PMS 515 PMS 516 PMS 517 PMS 518 PMS 519 PMS 520 PMS 521 PMS 522 PMS 523 PMS 524 PMS 525 PMS 526 PMS 527 PMS 528 PMS 529 PMS 530 PMS 531 PMS 532 PMS 533 PMS 534 PMS 535 PMS 536 PMS 537 PMS 538 PMS 539 PMS 540 PMS 541 PMS 542 PMS 543 PMS 544 PMS 545 PMS 546 PMS 547 PMS 548 PMS 549 PMS 550 PMS 551 PMS 552 PMS 553 PMS 554 PMS 555 PMS 556 PMS 557 PMS 558 PMS 559 PMS 560 PMS 561 PMS 562 PMS 563 PMS 564 PMS 565 PMS 566 PMS 567 PMS 568 PMS 569 PMS 570 PMS 571 PMS 572 PMS 573 PMS 574 PMS 575 PMS 576 PMS 577 PMS 578 PMS 579 PMS 580 PMS 581 PMS 582 PMS 583 PMS 584 PMS 585 PMS 586 PMS 587 PMS 588 PMS 589 PMS 590 PMS 591 PMS 592 PMS 593 PMS 594 PMS 595 PMS 596 PMS 597 PMS 598 PMS 599 PMS 600 PMS 601 PMS 602 PMS 603 PMS 604 PMS 605 PMS 606 PMS 607 PMS 608 PMS 609 PMS 610 PMS 611 PMS 612 PMS 613 PMS 614 PMS 615 PMS 616 PMS 617 PMS 618 PMS 619 PMS 620 PMS 621 PMS 622 PMS 623 PMS 624 PMS 625 PMS 626 PMS 627 PMS 628 PMS 629 PMS 630 PMS 631 PMS 632 PMS 633 PMS 634 PMS 635 PMS 636 PMS 637 PMS 638 PMS 639 PMS 640 PMS 641 PMS 642 PMS 643 PMS 644 PMS 645 PMS 646 PMS 647 PMS 648 PMS 649 PMS 650 PMS 651 PMS 652 PMS 653 PMS 654 PMS 655 PMS 656 PMS 657 PMS 658 PMS 659 PMS 660 PMS 661 PMS 662 PMS 663 PMS 664 PMS 665 PMS 666 PMS 667 PMS 668 PMS 669 PMS 670 PMS 671 PMS 672 PMS 673 PMS 674 PMS 675 PMS 676 PMS 677 PMS 678 PMS 679 PMS 680 PMS 681 PMS 682 PMS 683 PMS 684 PMS 685 PMS 686 PMS 687 PMS 688 PMS 689 PMS 690 PMS 691 PMS 692 PMS 693 PMS 694 PMS 695 PMS 696 PMS 697 PMS 698 PMS 699 PMS 700 PMS 701 PMS 702 PMS 703 PMS 704 PMS 705 PMS 706 PMS 707 PMS 708 PMS 709 PMS 710 PMS 711 PMS 712 PMS 713 PMS 714 PMS 715 PMS 716 PMS 717 PMS 718 PMS 719 PMS 720 PMS 721 PMS 722 PMS 723 PMS 724 PMS 725 PMS 726 PMS 727 PMS 728 PMS 729 PMS 730 PMS 731 PMS 732 PMS 733 PMS 734 PMS 735 PMS 736 PMS 737 PMS 738 PMS 739 PMS 740 PMS 741 PMS 742 PMS 743 PMS 744 PMS 745 PMS 746 PMS 747 PMS 748 PMS 749 PMS 750 PMS 751 PMS 752 PMS 753 PMS 754 PMS 755 PMS 756 PMS 757 PMS 758 PMS 759 PMS 760 PMS 761 PMS 762 PMS 763 PMS 764 PMS 765 PMS 766 PMS 767 PMS 768 PMS 769 PMS 770 PMS 771 PMS 772 PMS 773 PMS 774 PMS 775 PMS 776 PMS 777 PMS 778 PMS 779 PMS 780 PMS 781 PMS 782 PMS 783 PMS 784 PMS 785 PMS 786 PMS 787 PMS 788 PMS 789 PMS 790 PMS 791 PMS 792 PMS 793 PMS 794 PMS 795 PMS 796 PMS 797 PMS 798 PMS 799 PMS 800 PMS 801 PMS 802 PMS 803 PMS 804 PMS 805 PMS 806 PMS 807 PMS 808 PMS 809 PMS 810 PMS 811 PMS 812 PMS 813 PMS 814 PMS 815 PMS 816 PMS 817 PMS 818 PMS 819 PMS 820 PMS 821 PMS 822 PMS 823 PMS 824 PMS 825 PMS 826 PMS 827 PMS 828 PMS 829 PMS 830 PMS 831 PMS 832 PMS 833 PMS 834 PMS 835 PMS 836 PMS 837 PMS 838 PMS 839 PMS 840 PMS 841 PMS 842 PMS 843 PMS 844 PMS 845 PMS 846 PMS 847 PMS 848 PMS 849 PMS 850 PMS 851 PMS 852 PMS 853 PMS 854 PMS 855 PMS 856 PMS 857 PMS 858 PMS 859 PMS 860 PMS 861 PMS 862 PMS 863 PMS 864 PMS 865 PMS 866 PMS 867 PMS 868 PMS 869 PMS 870 PMS 871 PMS 872 PMS 873 PMS 874 PMS 875 PMS 876 PMS 877 PMS 878 PMS 879 PMS 880 PMS 881 PMS 882 PMS 883 PMS 884 PMS 885 PMS 886 PMS 887 PMS 888 PMS 889 PMS 890 PMS 891 PMS 892 PMS 893 PMS 894 PMS 895 PMS 896 PMS 897 PMS 898 PMS 899 PMS 900 PMS 901 PMS 902 PMS 903 PMS 904 PMS 905 PMS 906 PMS 907 PMS 908 PMS 909 PMS 910 PMS 911 PMS 912 PMS 913 PMS 914 PMS 915 PMS 916 PMS 917 PMS 918 PMS 919 PMS 920 PMS 921 PMS 922 PMS 923 PMS 924 PMS 925 PMS 926 PMS 927 PMS 928 PMS 929 PMS 930 PMS 931 PMS 932 PMS 933 PMS 934 PMS 935 PMS 936 PMS 937 PMS 938 PMS 939 PMS 940 PMS 941 PMS 942 PMS 943 PMS 944 PMS 945 PMS 946 PMS 947 PMS 948 PMS 949 PMS 950 PMS 951 PMS 952 PMS 953 PMS 954 PMS 955 PMS 956 PMS 957 PMS 958 PMS 959 PMS 960 PMS 961 PMS 962 PMS 963 PMS 964 PMS 965 PMS 966 PMS 967 PMS 968 PMS 969 PMS 970 PMS 971 PMS 972 PMS 973 PMS 974 PMS 975 PMS 976 PMS 977 PMS 978 PMS 979 PMS 980 PMS 981 PMS 982 PMS 983 PMS 984 PMS 985 PMS 986 PMS 987 PMS 988 PMS 989 PMS 990 PMS 991 PMS 992 PMS 993 PMS 994 PMS 995 PMS 996 PMS 997 PMS 998 PMS 999 PMS 1000	<p>MDC 0088-1799-42</p> <p>Cardizem® CD diltiazem HCl</p> <p>CD 360mg</p> <p>ONCE-A-DAY DOSAGE</p> <p>90 Capsules Hoechst Marion Roussel</p>	<p>Each CARDIZEM® CD capsule contains 360 mg of diltiazem hydrochloride. ONLY Dosage and Administration Read package insert for prescribing information. WARNING: Keep out of reach of children. Pharmacist: Dispense in light-resistant, tight container with child-resistant closure. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) (see USP Controlled Room Temperature). Avoid excessive humidity.</p> <p>©1998 Hoechst Marion Roussel, Inc. Hoechst Marion Roussel, Inc. Kansas City, MO 64137 USA www.jrn.com</p>	<p>3 0088-1799-42 1</p> <p>50018907</p>
--	---	---	---

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

CHEMISTRY REVIEW(S)

JUL 28 1999

CHEMIST'S REVIEW		1. ORGANIZATION HFD-110	2. NDA Number 20-062
3. Name and Address of Applicant (City & State) Carderm Capital L.P. c/o Westbroke Limited Raymond House 12 Par La Ville Road Hamilton, HM 12 Bermuda		4. Supplement(s) Number(s) Date(s) SCP-027 6/18/99	
5. Drug Name CARDIZEM CD	6. Nonproprietary Name Diltiazem hydrochloride		8. Amendments & Other (reports, etc) - Dates Orig - 1/7/99 BC-3/5/99
7. Supplement Provides For: Response to letter of May 7, 1999 for S-027.			
9. Pharmacological Category Ca antagonist (hypertension)	10. How Dispensed <input checked="" type="checkbox"/> Rx <input type="checkbox"/> OTC		11. Related IND(s)/NDA(s)/DMP(s)
12. Dosage Form(s) Capsules, CD (controlled diffusion, once-a-day)	13. Potency(ies) 120, 180, 240 and 300 mg/capsule		
14. Chemical Name and Structure 1,5-Benzothiazepin-4(5H)one, 3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2-(4-(dimethylphenyl)-, monohydrochloride, (+)-cis-		15. Records/Reports Current <input type="checkbox"/> Yes <input type="checkbox"/> No Reviewed <input type="checkbox"/> Yes <input type="checkbox"/> No	
16. Comments: As requested in the letter, final proofs for the bottle labels and final printed labeling for the package insert showing the changes to the storage statement are included. Updated (12 month) stability report is included. The statistical analysis shows that the projected shelf life of the 360 mg capsules is greater than 24 months is also included. Labels - storage statement has been corrected. Satisfactory.			
17. Conclusions and Recommendations: For the final printed labeling, Hoechst Marion Roussel needs to use larger font. Unacceptable.			
18. REVIEWER			
Name Danute G. Cunningham	Signature 		Date Completed June 29, 1999
Distribution: <input checked="" type="checkbox"/> Original Jacket <input type="checkbox"/> Reviewer <input type="checkbox"/> Division File <input type="checkbox"/> CSO			

20062S27.AM1

1348-99

MAY 6 1999

CHEMIST'S REVIEW		1. ORGANIZATION HFD-110		2. NDA Number 20-062	
3. Name and Address of Applicant (City & State) Carderm Capital L.P. c/o Westbroke Limited Raymond House 12 Par La Ville Road Hamilton, HM 12 Bermuda				4. Supplement(s) Number(s) Date(s) SCF-027 1/7/99	
5. Drug Name CARDIZEM CD		6. Nonproprietary Name Diltiazem hydrochloride		8. Amendments & Other (reports, etc) - Dates B: 5/5/99 etc	
7. Supplement Provides For: strength addition of 360 mg capsule (slightly modified) to the Cardizem CD capsules.					
9. Pharmacological Category Ca antagonist (hypertension)		10. How Dispensed <input checked="" type="checkbox"/> Rx <input type="checkbox"/> OTC		11. Related IND(s)/NDA(s)/DMF(s) B	
12. Dosage Form(s) Capsules, CD (controlled diffusion, once-a-day)		13. Potency(ies) 120, 180, 240 and 300 mg/capsule			
14. Chemical Name and Structure 1,5-Benzothiazepin-4(5H)one, 3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2-(4-(dimethylphenyl)-, monohydrochloride, (+)-cis-				15. Records/Reports Current <input type="checkbox"/> Yes <input type="checkbox"/> No Reviewed <input type="checkbox"/> Yes <input type="checkbox"/> No	
16. Comments: The formulation change is found in the Active Bead. All other aspects, including the sustained release coating, of the manufacturing process of the capsule remain unchanged from that which is currently approved. This new capsule will be manufactured and controlled by Hoechst Marion Roussel, Inc. in Kansas City, MO. No change in formulation to the other approved strengths is proposed in this supplement.					
17. Conclusions and Recommendations: EES requested on 1/19/99. Acceptable on 5/5/99 Biopharmaceutics review requested on 1/19/99. Approvable - 4/23/99. Approvable - due to labeling issues. Storage statement has to be modified: Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Container label - storage statement should be changed.					
18. REVIEWER					
Name Danute G. Cunningham		Signature <i>DS</i>		Date Completed May 5, 1999	
Distribution: <input checked="" type="checkbox"/> Original Jacket <input type="checkbox"/> Reviewer <input type="checkbox"/> Division File <input type="checkbox"/> CSO					

20062S27.SUP

5/5/99

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

**CLINICAL PHARMACOLOGY AND
BIOPHARMACEUTICS REVIEW(S)**

APR 23 1999

only

Clinical Pharmacology/Biopharmaceutics Review

NDA 20-062

Serial #: SCF-027

Compound #: Cardizem CD 360mg Capsules

Hoechst Marion Roussel

Submission Date: January 7, 1999

Reviewer: Thomas A. Parmelee, Pharm.D.

Type of Submission: Supplement for New Formulation Study Report-Cardizem CD 360mg Capsules- A Bioequivalence Study and a Food Effect Study

BACKGROUND

NDA-062 has been approved for Cardizem CD (diltiazem HCl) Capsules in the strengths of 120mg, 180mg, 240mg, and 300mg. The maximum daily dose for diltiazem extended release capsules is established at 360mg. A new 360mg capsule formulation has been developed, and is the topic of this supplemental submission. The new 360mg capsule contains a formulation that is slightly modified from the currently approved lower strength capsules. The formulation change is found in the active bead of the drug product.

Two studies were submitted to the Office of Clinical Pharmacology and Biopharmaceutics for review. These studies were designed to show that the new Cardizem CD 360mg capsule formulation is bioequivalent to two Cardizem CD 180mg marketed capsules. One study is a bioequivalence study comparing single-dose and multiple-dose administration of the new 360mg capsules to the marketed 180mg strength capsules. The second study examines the effect of food on the single-dose pharmacokinetics of the new 360mg diltiazem capsule formulation. These studies are summarized in Appendix 1 and Appendix 2, respectively.

RESULTS

It appears that both lots of the new 360mg capsule formulation are bioequivalent to the marketed 180mg capsule formulation in the single-dose comparisons in terms of both AUC (0-inf) and C_{max} for both parent diltiazem and N-desmethyl metabolite.

In the steady-state comparisons, bioequivalency is met in terms of AUC, ss and C_{max}, ss between treatments, and only fails the 80-125% rule for Treatment B (lot # RH9738) in terms of C_{min}, ss for parent diltiazem.

The 90% confidence intervals between the high-fat breakfast treatment and the fasted treatment were within the 80-125% rule for both AUC (0-inf) and Cmax when looking at parent diltiazem and the N-desmethyl metabolite. Food does not appear to significantly affect the PK parameters of either lot of 360mg diltiazem capsules.

COMMENTS

- 1) Gender should not have been considered inclusion/exclusion criteria for these two clinical studies unless there was a specific reason for doing so. This point was mentioned to the sponsor via a teleconference before the start of the study.
- 2) The dissolution specifications are appropriate for the new strength capsule.
- 3) The draft prescription labeling submitted from the sponsor shows that the 360mg capsules contain black iron oxide, FD&C Blue #1, and starch. These ingredients were not listed in the composition of the capsules for review.

RECOMMENDATIONS

The new dosage strength for Cardizem CD 360mg capsules is approvable from the standpoint of the Office of Clinical Pharmacology and Biopharmaceutics. The comment above regarding the draft prescription labeling was conveyed to the review chemist. The dissolution methodology and specifications for the new strength are:

Apparatus: USP Type 2 (paddle)
Speed: 100 rpm
Media: 900mL degassed 0.1N HCl
Temperature: 37 C +/- 0.5 C

<u>Time (hrs.)</u>	<u>Specifications (%)</u>
6 hours	%
12 hours	%
18 hours	%
24 hours	NLT %
30 hours	NLT %

The draft prescription labeling (updated October 1998) and label included with the submission are attached to this review. The labeling for all diltiazem products is currently being updated and reviewed by this division (updated November 1998). The labeling for this new Cardizem CD 360mg capsule formulation should reflect the final printed labeling decided upon by the sponsor and the Agency for all Cardizem CD products.

IS/
Thomas A. Parmelee, Pharm.D.

4/23/99

APPENDIX 1

"BIOEQUIVALENCE OF 360MG DILTIAZEM HCL FORMULATIONS AND CARDIZEM CD AFTER SINGLE AND MULTIPLE DOSE ADMINISTRATIONS IN HEALTHY MALE SUBJECTS"

STUDY: Protocol # DZPR0207
Report K-98-0235-D

SPONSOR: Licensed to:
Hoechst Marion Roussel Inc.
P.O. Box 9627, H3-M2112
Kansas City, MO 64134-0627

Authorized by:
Carderm Capital L.P.
Raymond House
12 Par La Ville Road
Hamilton, HM 12 Bermuda

INVESTIGATOR AND STUDY SITE:

OBJECTIVES:

To determine whether 360mg Diltiazem HCL capsule formulations are bioequivalent to marketed 180mg Cardizem CD capsules.

FORMULATIONS:

- 1) Diltiazem 360mg capsules (lot# RH9736); Batch size
- 2) Diltiazem 360mg capsules (lot# RH9738); Batch size
- 3) Cardizem CD 180mg marketed capsules (lot# P31048)

The following table shows the composition of the new formulation of Cardizem CD 360mg Capsules:

STUDY DESIGN:

The study design was a randomized, open-label, single- and multiple-dose, 3-period, crossover study with a washout period of 12 days between treatments. The study population was 26 healthy, non-smoking males between the ages of 18 to 45 years. Subjects received each of the three treatment regimens in a randomized fashion:

Treatment A: One diltiazem 360mg capsule (RH9736) given as a SD on day 1, and then q.d. on days 3-9.

Treatment B: One diltiazem 360mg capsule (RH9738) given as a SD on day 1, and then q.d. on days 3-9.

Treatment C: Two Cardizem CD 180mg capsules (P31048) given as a SD on day 1, and then q.d. on days 3-9.

Subjects were continuously monitored for general health and any adverse reactions. Heart rate, blood pressure, and 12-lead ECG recordings, clinical chemistry, and hematological exams were done before the study and upon completion. Plasma samples were collected before the SD on day 1 and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, 24, 36, and 48 hours following the dose. The subjects received seven days of multiple dosing during days 3-9. Trough plasma samples were obtained before the dose on days 8 and 9, and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, and 24 hours following the day 9 dose.

ASSAY:

Table B: Dissolution Data for Cardizem CD 360mg Capsules

Time	Specifications (%)	RH9738 (%)	Percent Dissolved RH9736 (%)	P31048 180mg (%)
3 hours				
6 hours	%			
9 hours				
12 hours	%			
15 hours				
18 hours	%			
24 hours	NLT %			
30 hours	NLT %			

DATA ANALYSIS:

Pharmacokinetic analysis of diltiazem and MA metabolite concentrations in plasma was conducted by non-compartmental methods. The metabolite DAD concentrations were presented by descriptive statistics only (mean, standard deviation, CV%). The primary PK comparisons include C_{max} and AUC (0-inf) following single dose administration; and C_{max,ss}, C_{min,ss}, trough plasma concentrations (days 8, 9, and 10), and AUC_{ss} for multiple dose steady-state findings.

Comparisons between treatments were made for diltiazem and MA metabolite pharmacokinetic parameters and trough plasma levels. An analysis of variance (ANOVA) was performed for each parameter using PROC MIXED SAS with terms for sequence, subject within sequence, period, and treatment. Least square means, treatment differences, and 90% confidence intervals for treatment differences were determined. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, treatment ratios, and 90% confidence intervals for these treatment ratios. Each lot of the diltiazem 360mg (Treatments A and B) was compared to the marketed reference Cardizem CD 180mg (Treatment C). Bioequivalence was to

be concluded if the limits of the 90% confidence interval on the ratio of treatment means falls entirely within the 80-125% range.

Trough plasma concentrations for each treatment were also compared using an ANOVA with terms for subject and day. From this ANOVA, least square means for each day, estimated differences between days, and 90% confidence intervals for the differences between days were calculated. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, day ratios, and 90% confidence intervals for these ratios.

RESULTS:

Both lots of the 360mg capsules appear to be bioequivalent to the reference Cardizem CD 180mg capsules in the single-dose comparisons. Treatment A (lot # RH9736) appears to be bioequivalent to the reference capsules in the multiple-dose steady state comparison, however, treatment B (lot # 9738) is outside the 80-125% BE limits for C_{min}, ss. Please refer to the following tables and figures:



Figure 3. Mean diltiazem plasma concentrations following 360 mg single dose of once-daily capsules on day 1; protocol D2P10207

A - lot # RH9736 based (n=24); B - lot # 9738 based (n=23); C - lot # P1743 based (n=23); Cardizem CD




Figure 4. Mean diltiazem plasma concentrations following 360 mg dose (time 0) of once-daily capsules on day 5; -24 hours = trough sample on day 5; Protocol D2P10207

A - lot # RH9736 based (n=24); B - lot # 9738 based (n=23); C - lot # P1743 based (n=23); Cardizem CD




Figure 5. Mean N-desmethylidiazepam plasma concentrations following 360 mg single dose on day 1 of once-daily capsules. Protocol DZPR0207

A: 12 P10736 tested (n=24); B: 12 P10736 tested (n=24); C: 12 P10736 tested (n=24)
(Continued on next page)




Figure 6. Mean N-desmethylidiazepam plasma concentrations following 360 mg dose (time 0) on day 2 of once-daily capsules. 24 hours = trough sample on day 2. Protocol DZPR0207

A: 12 P10736 tested (n=24); B: 12 P10736 tested (n=24); C: 12 P10736 tested (n=24)
(Continued on next page)

30 October FINAL

Table 9. Mean diltiazem (DTZ) pharmacokinetic parameters following 360 mg single dose on day 1, Protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC (0-∞) (ng/mL·h)	A	24	3437.58	3254.72	30.91	A/C	100.83	(88.5, 114.8)	0.916
	B	23	3676.08	3436.67	36.23	B/A	105.59	(92.7, 120.3)	0.487
	C	23	3478.85	3228.07	39.54	B/C	106.46	(93.4, 121.3)	0.425
C _{max} (ng/mL)	A	24	170.69	160.18	38.54	A/C	102.47	(90.6, 115.8)	0.740
	B	23	169.69	158.35	35.20	B/A	98.86	(87.4, 111.8)	0.876
	C	23	166.58	156.32	36.49	B/C	101.30	(89.6, 114.5)	0.861
t _{1/2} (h)	A	24	6.98	6.86	16.61	A/C	95.33	(86.7, 104.8)	0.403
	B	23	7.48	7.10	45.65	B/A	103.51	(94.1, 113.9)	0.546
	C	23	7.30	7.20	19.48	B/C	98.68	(89.7, 108.6)	0.816
t _{max} (h)	A	24	13.08	12.17	35.13	A/C	106.27	(88.0, 128.3)	0.589
	B	23	13.22	12.45	30.52	B/A	102.34	(84.8, 123.6)	0.837
	C	23	12.39	11.45	34.30	B/C	108.77	(90.0, 131.5)	0.460

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given fasted

TRT B = one 160 mg CD capsule (lot RH9738) given fasted

TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given fasted

Supporting Data:

Appendix B.3.3 Details of treatment comparisons, diltiazem single dose pharmacokinetic parameters, page 214 and Appendix C.2.2 Pharmacokinetic listings, page 639

30 October FINAL

Table 10. Mean diltiazem (DTZ) steady-state pharmacokinetic parameters following 360 mg dose on day 9, protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^b on ratio	P value
AUC _{ss} (ng/mLxh)	A	24	3754.53	3551.98	28.57	A/C	100.69	(94.0, 107.9)	0.868
	B	23	3896.27	3558.25	36.19	B/A	100.18	(93.5, 107.3)	0.966
	C	23	3811.93	3527.75	33.13	B/C	100.86	(94.2, 108.1)	0.834
C _{max,ss} (ng/mL)	A	24	224.18	212.39	29.46	A/C	89.55	(83.5, 96.1)	0.011
	B	23	245.21	225.41	34.48	B/A	106.13	(98.9, 113.9)	0.163
	C	23	256.14	237.17	33.81	B/C	95.04	(88.6, 101.9)	0.230
C _{min,ss} (ng/mL)	A	24	97.29	87.97	38.53	A/C	104.02	(92.8, 116.6)	0.564
	B	23	109.15	97.94	41.76	B/A	111.33	(99.3, 124.8)	0.122
	C	23	94.05	84.57	41.10	B/C	115.80	(103.3, 129.8)	0.036
RATIO (C _{max,ss} / C _{min,ss})	A	24	2.55	2.41	49.16	A/C	85.77	(77.1, 95.4)	0.020
	B	23	2.36	2.31	25.23	B/A	95.78	(86.1, 106.6)	0.501
	C	23	2.89	2.81	29.11	B/C	82.14	(73.8, 91.4)	0.004
t _{max} (h)	A	24	9.75	9.38	43.75	A/C	149.54	(127.7, 175.1)	<0.001
	B	23	6.65	6.30	35.86	B/A	67.15	(57.3, 78.7)	<0.001
	C	23	6.65	6.27	36.15	B/C	100.42	(85.8, 117.5)	0.965
Trough Plasma Conc ^a (ng/mL)	A	24	116.96	110.78	31.37	A/C	108.12	(100.8, 116.0)	0.069
	B	23	119.28	110.00	36.39	B/A	99.29	(92.5, 106.5)	0.866
	C	23	110.28	102.46	34.20	B/C	107.36	(100.1, 115.2)	0.097

^a mean of trough plasma concentrations on days 8, 9, and 10

^b percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given on days 3 through 9

TRT B = one 360 mg CD capsule (lot RH9738) given on days 3 through 9

TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given on days 3 through 9

Supporting Data:

Appendix B.3.7 Details of treatment comparisons, diltiazem steady state pharmacokinetic parameters, page 223

Appendix C.2.2 Pharmacokinetic listings, page 639

Table 11. Mean N-desmethyldiltiazem (MA) pharmacokinetic parameters following 360 mg dose on day 1, Protocol DZPR0207									
	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC(0-∞) (ng/mLxh)	A	24	1246.89	1176.76	32.17	A/C	99.12	(87.2, 112.6)	0.907
	B	23	1402.22	1272.85	56.64	B/A	108.17	(95.2, 122.9)	0.309
	C	23	1263.94	1187.27	36.93	B/C	107.21	(94.3, 121.9)	0.866
C _{max} (ng/mL)	A	24	43.05	40.43	35.32	A/C	97.49	(87.5, 108.6)	0.695
	B	23	43.37	41.12	28.05	B/A	101.72	(91.3, 113.4)	0.792
	C	23	43.86	41.47	29.52	B/C	99.17	(89.0, 110.5)	0.898
T _{1/2} (h)	A	24	11.16	10.96	18.53	A/C	99.14	(86.2, 114.0)	0.917
	B	23	13.79	12.09	86.90	B/A	110.31	(95.9, 126.9)	0.245
	C	23	11.22	11.06	23.65	B/C	109.36	(95.0, 125.9)	0.291
T _{max} (h)	A	24	16.63	16.12	22.49	A/C	100.83	(88.2, 115.2)	0.918
	B	23	16.52	15.38	47.86	B/A	95.42	(83.5, 109.1)	0.559
	C	23	16.09	15.99	19.20	B/C	96.21	(84.1, 110.0)	0.631
^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data TRT A = one 360 mg CD capsule (lot RH9736) given fasted TRT B = one 360 mg CD capsule (lot RH9738) given fasted TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given fasted Supporting Data: Appendix B.3.14 Details of treatment comparisons, MA single dose pharmacokinetic parameters, page 237 Appendix C.2.2 Pharmacokinetic listings, page 639									

30 October FINAL

Table 12. Mean N-desmethyldiltiazem (MA) steady-state pharmacokinetic parameters following 360 mg dose on day 8, Protocol DZPR0207

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^b on ratio	P value
ADC _{ss} (ng/mLxh)	A	24	1333.87	1254.56	31.56	A/C	98.30	(94.1, 102.7)	0.512
	B	23	1344.84	1254.89	33.11	B/A	100.03	(95.7, 104.5)	0.992
	C	23	1365.51	1276.27	33.37	B/C	98.33	(94.1, 102.7)	0.517
C _{max,ss} (ng/mL)	A	24	70.41	66.52	31.41	A/C	97.52	(93.0, 102.2)	0.376
	B	23	68.45	64.15	30.65	B/A	96.45	(92.0, 101.1)	0.205
	C	23	72.68	68.21	32.45	B/C	94.05	(89.7, 98.6)	0.034
C _{min,ss} (ng/mL)	A	24	41.31	37.48	39.71	A/C	102.20	(95.1, 109.8)	0.614
	B	23	43.80	40.62	35.07	B/A	108.36	(100.8, 116.5)	0.068
	C	23	40.07	36.68	38.79	B/C	110.73	(103.1, 119.0)	0.022
RATIO (C _{max,ss} / C _{min,ss})	A	24	1.82	1.78	33.29	A/C	95.41	(89.5, 101.8)	0.227
	B	23	1.59	1.58	12.60	B/A	89.04	(83.5, 95.0)	0.004
	C	23	1.88	1.86	20.76	B/C	84.95	(79.6, 90.6)	<0.001
t _{max} (h)	A	24	13.54	11.90	31.40	A/C	115.57	(92.6, 144.2)	0.278
	B	23	10.00	9.11	43.06	B/A	76.55	(61.3, 95.6)	0.049
	C	23	11.26	10.30	36.73	B/C	88.47	(70.8, 110.6)	0.361
Trough Plasma Conc ^a (ng/mL)	A	24	47.21	44.67	32.31	A/C	105.95	(101.4, 110.7)	0.032
	B	23	46.66	43.85	33.66	B/A	98.18	(94.0, 102.6)	0.486
	C	23	44.84	42.16	33.08	B/C	104.02	(99.6, 108.7)	0.137

^a mean of trough plasma concentrations on days 8, 9, and 10
^b percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data
 TRT A = one 360 mg CD capsule (lot RH9736) given on days 3 through 9
 TRT B = one 360 mg CD capsule (lot RH9738) given on days 3 through 9
 TRT C = two 180 mg Cardizem CD Capsule (lot P31048) given on days 3 through 9
 Supporting Data:
 Appendix B.3.18 Details of treatment comparisons, MA steady state pharmacokinetic parameters, page 246 and Appendix C.2.2 Pharmacokinetic listings, page 639

APPENDIX 2

"EFFECT OF FOOD ON THE SINGLE-DOSE PHARMACOKINETICS OF DILTIAZEM HCl 360MG FORMULATIONS IN HEALTHY MALE SUBJECTS"

STUDY: Protocol # DZPR0208
Report K-98-0236-D

SPONSOR: Licensed to:
Hoechst Marion Roussel Inc.
P.O. Box 9627, H3-M2112
Kansas City, MO 64134-0627

Authorized by:
Carderm Capital L.P.
Raymond House
12 Par La Ville Road
Hamilton, HM 12 Bermuda

INVESTIGATOR AND STUDY SITE:

OBJECTIVES:

To determine the effects of a high-fat breakfast on the rate and extent of absorption of a single oral dose of 360mg diltiazem HCl capsule formulation.

FORMULATIONS:

- 1) Diltiazem HCl 360mg capsules (lot# RH9736)
- 2) Diltiazem HCl 360mg capsules (lot# RH9738)

STUDY DESIGN:

The study design was a randomized, open-label, single-dose 4-period, crossover study with a washout period of 7 days between treatments. The study population was 22 healthy, non-smoking males between the ages of 18 to 45 years. Subjects received each of the four treatment regimens in a randomized fashion:

Treatment A: One diltiazem 360mg capsule (RH9736) dosed under fasting conditions.

Treatment B: One diltiazem 360mg capsule (RH9736) dosed with a high-fat breakfast.
Treatment C: One diltiazem 360mg capsule (RH9738) dosed under fasting conditions.
Treatment D: One diltiazem 360mg capsule (RH9738) dosed with a high-fat breakfast.

Subjects were continuously monitored for general health and adverse events. Heart rate, blood pressure, and 12-lead ECG recordings, clinical chemistry, and hematological exams were done before the study and upon completion. Heart rate, blood pressure (5 minutes supine), and lead II ECG measurements were taken 4 hours following each single dose. Plasma samples were collected before each dose on day 1 and 1, 2, 3, 4, 5, 6, 8, 10, 12, 14, 16, 18, 21, 24, 36, and 48 hours following the dose.

ASSAY:

DATA ANALYSIS:

Pharmacokinetic analysis of diltiazem and MA metabolite concentrations in plasma was conducted by non-compartmental methods. The metabolite DAD concentrations were presented by descriptive statistics only (mean, standard deviation, CV%). The primary PK comparisons include C_{max} and AUC (0-inf) for plasma concentrations.

Comparisons between treatments were made for diltiazem and MA metabolite pharmacokinetic parameters. An analysis of variance (ANOVA) was performed for each parameter using PROC MIXED SAS with terms for sequence, subject within sequence, period, and treatment. Least square means, treatment differences, and 90% confidence intervals for treatment differences were determined. These log-transformed results were back-transformed by exponentiation to obtain adjusted means, treatment ratios, and 90% confidence intervals for these treatment ratios. Treatment B was compared to Treatment A with Treatment A serving as the reference, and Treatment D was compared to Treatment C with Treatment C as the reference treatment. Equivalence was defined as the limits of the 90% confidence interval on the ratio of treatment means falling entirely within 80% to 125%.

RESULTS:

Twenty subjects completed all four treatments. The differences in AUC (0-inf) and Cmax between the high-fat and fasting treatments were small. The 90% confidence intervals for the differences between treatments were within the limits of 80% to 125% using the fasted treatments as the references. Please refer to the following tables and figures:

**APPEARS THIS WAY
ON ORIGINAL**




Figure 3. Mean diltiazem plasma concentrations following 360 mg single dose of once-daily capsules, Protocol DZPR0208

A= lot RH9736 fasted (n=20), B= lot RH9736 fed (n=20), C= lot RH9738 fasted (n=20), D= lot RH9738 fed (n=21).

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings,

page 473




Figure 4. Mean N-desmethyldiltiazem plasma concentrations following 360 mg single dose of once-daily capsules, Protocol DZPR0208

A= lot RH9736 fasted (n=20), B= lot RH9736 fed (n=20), C= lot RH9738 fasted (n=20), D= lot RH9738 fed (n=21).

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings,

page 473




Figure 5. Mean DAD plasma concentrations, Protocol DZPR0208

A= lot RH9736 fasted, B= lot RH9736 fed, C= lot RH9738 fasted, D= lot RH9738 fed.

Supporting Data:
Appendix C.2.2 Pharmacokinetic listings,

page 473

Table 9. Mean diltiazem (DTZ) pharmacokinetic parameters, 360 mg single dose, protocol DZPR0208

	TR T	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC(0—) (ng/mL·h)	A	20	3384.33	3106.43	29.03	—	—	—	—
	B	20	3517.62	3240.02	31.23	B/A	104.30	(95.1, 114.4)	0.451
	C	20	3214.98	2961.00	27.04	—	—	—	—
	D	21	3633.28	3272.02	47.82	D/C	110.50	(100.7, 121.2)	0.077
C _{max} (ng/mL)	A	20	160.48	149.61	30.10	—	—	—	—
	B	20	179.60	166.93	34.52	B/A	111.58	(101.8, 122.3)	0.051
	C	20	153.51	144.68	24.63	—	—	—	—
	D	21	174.18	159.05	43.74	D/C	109.93	(100.3, 120.5)	0.069
t _{1/2} (h)	A	20	6.87	6.68	16.06	—	—	—	—
	B	20	6.65	6.49	13.47	B/A	97.16	(92.7, 101.8)	0.306
	C	20	6.77	6.60	13.55	—	—	—	—
	D	21	6.49	6.41	16.21	D/C	97.03	(92.6, 101.7)	0.283
t _{max} (h)	A	20	11.40	10.15	45.93	—	—	—	—
	B	20	10.10	9.33	40.37	B/A	91.93	(73.3, 115.3)	0.536
	C	20	13.00	11.85	38.18	—	—	—	—
	D	21	10.48	9.21	57.24	D/C	77.73	(62.2, 97.2)	0.065

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data

TRT A = one 360 mg CD capsule (lot RH9736) given fasted

TRT B = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

TRT C = one 360 mg CD capsule (lot RH9736) given fasted

TRT D = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

Supporting Data:

Appendix B.3.3 Details of treatment comparisons, page 201 and Appendix C.2.2 Pharmacokinetic listings, page 473

Table 10. Mean N-desmethyldiltiazem (MA) pharmacokinetic parameters, 360 mg single dose, protocol DZPR0208

	TRT	Number	Raw mean	Adjusted mean	CV%	Pair	Ratio (%)	90% CI ^a on ratio	P value
AUC (0--)(ng/mL·h)	A	20	1161.61	1083.07	27.09	B/A	104.62	(97.7, 112.0)	0.272
	B	20	1196.27	1133.09	23.05				
	C	20	1081.72	1011.27	24.71				
	D	21	1166.22	1116.67	29.43				
C _{max} (ng/mL)	A	20	41.29	39.18	27.14	B/A	106.38	(99.8, 113.4)	0.109
	B	20	43.22	41.68	25.04				
	C	20	38.12	36.43	21.19				
	D	21	40.04	38.85	24.78				
t _{1/2} (h)	A	20	10.22	9.89	17.45	B/A	101.20	(96.3, 106.4)	0.689
	B	20	10.32	10.01	17.03				
	C	20	9.97	9.64	17.05				
	D	21	10.40	10.15	22.52				
t _{max} (h)	A	20	16.85	16.29	18.99	B/A	70.88	(58.6, 85.7)	0.004
	B	20	12.45	11.55	35.97				
	C	20	16.10	15.77	18.68				
	D	21	12.14	10.63	50.82				
						D/C	68.68	(56.9, 82.9)	0.001

^a percent ratio and 90% confidence interval (CI) were calculated from ANOVA using log transformed data
 TRT A = one 360 mg CD capsule (lot RH9736) given fasted
 TRT B = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast
 TRT C = one 360 mg CD capsule (lot RH9736) given fasted
 TRT D = one 360 mg CD capsule (lot RH9736) given with high-fat breakfast

Supporting Data:

Appendix B.3.7 Details of treatment comparisons, page 209 and Appendix C.2.2 Pharmacokinetic listings, page 473

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

ADMINISTRATIVE DOCUMENTS

RHPM Review of Final Printed Labeling

AUG 24 1999

Application: NDA 20-062
Cardizem CD (diltiazem HCl) Capsules

Applicant: Carderm Capital L.P.

Supplement Date: January 7, 1999

FPL Letter Date: June 18, 1999

FPL Receipt Date: June 21, 1999

Background

NDA 20-062/S-027 provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules. An approvable letter was issued on May 7, 1999. In addition to the labeling changes under **DESCRIPTION** and **HOW SUPPLIED** relating to the new dosage strength, the approvable letter requested a revision of the **Storage Statement**.

Review

The applicant submitted final printed labeling in a submission dated June 18, 1999. The labeling was revised to include information on the 360 mg capsule under **DESCRIPTION** and **HOW SUPPLIED**. In addition, the **Storage Statement** was revised to read as follows:

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

These changes were made in accordance with the requests in the approvable letter. An approval letter will be drafted for Dr. Lipicky's signature.


David Roeder
Regulatory Health Project Manager

cc: NDA 20-062
HFD-110
HFD-110/DRoeder/ABlount

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER: NDA 20062/S027

CORRESPONDENCE

JUL 19 1999

Hoechst Marion Roussel, Inc.
Attention: Janet K. DeLeon
10236 Marion Park Drive
P.O. Box 9627
Kansas City, MO 64134-0627

Dear Ms. DeLeon:

Please refer to your January 7, 1999 supplemental new drug application (NDA) submitted under section 505(b) of the Federal Food, Drug and Cosmetic Act for CARDIZEM CD (diltiazem hydrochloride) Capsules, 180 mg, 240 mg, 300 mg and 360 mg.

The supplemental application provides for a new dosage strength, 360 mg Capsules. The formulation of this new capsule strength is slightly modified from the other approved dosage strength capsules.

We have completed our validation of the analytical methods for the 360 mg capsules and request the following additional information regarding the dissolution test:

The method refers to dissolution software used to correct for UV absorbance interference from diethyl phthalate, an excipient in the product. Attempts on April 8, 1999 by the analyst to get detailed information and explicit calculation formulas from your firm for dissolution calculations for the excipient contribution were not entirely successful. Please include a detailed description of the software and the calculations used to obtain the final results in the method.

The method does not specify whether aliquots taken out are replaced or not. If not replaced, please state whether final results are corrected for the volume taken during sampling. The validating analyst did not replace aliquots and corrected the volume withdrawn. It may be that sample aliquots are circulated back into the dissolution bath after samples are read. If this is the case, it should be stated in the method.

We would appreciate your prompt written response.

If you have any questions, please contact Danute G. Cunningham at (301) 594-5351 or Kasturi Srinivasachar, Ph.D. at (301) 594-5376.

Sincerely yours,

JSI 7-19-99
Kasturi Srinivasachar, Ph.D.
Chemistry Team Leader, DNDC I, for the
Division of Cardio-Renal Drug Products, (HFD-110)
Office of New Drug Chemistry
Center for Drug Evaluation and Research